

## COLORECTAL BUNDLE CHECKLIST ON PARAGON

AIN	INF SCRIN	HISTORY 1	HISTORY 2	REVIEW	VAL INFO	EENT	NEURO	RESP	CARDIO	GASTRO	GENITC
-----	-----------	-----------	-----------	--------	----------	------	-------	------	--------	--------	--------

### Pre-operative (ASC)

- Give patient the SSI patient education sheet
- Encourage smoking cessation 30 days before surgery
- Use preoperative skin cleansing: with chlorhexidine wipes (night before and morning of surgery)
- Mechanically prepare the colon the day before surgery
- Administer non-absorbable oral antimicrobial agents (neomycin and metronidazole) the night before surgery
- Screen diabetic and non-diabetic patients using HbA1c levels
- Check blood glucose levels, if >140mg/dl, start insulin infusion
- Remove hair with clippers in holding area
- Apply forced warm air gown to maintain normothermia
- Other

### Intraoperative (OR)

- Prescribe appropriate antibiotic
- Dose prophylactic antimicrobial agent based on weight
- Administer prophylactic antimicrobial agents IV on time
- Redose prophylactic antibiotic based on duration of operation
- Use standardized antiseptic agent for skin preparation: chlorhexidine gluconate (Chloraprep)
- Use at least 80% fraction of inspired oxygen
- Ensure double gloving for all scrubbed surgical team members
- Maintain perioperative normothermia
- Aggressively control glucose in all patients, start insulin infusions for any blood glucose level >140mg/dl
- Other

### Postoperative (PACU)

- Maintain control of serum blood glucose levels in all patients, glycemic control team (anesthesia) consulted
- Protect primary incisions with antimicrobial dressing for 5 days
- Use high fraction of inspired oxygen (non-rebreather mask) for 4 hours
- Discontinue prophylactic antimicrobial agent within 24 hours of surgery
- Other